

U

DTIC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE

## DTIC REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION

ELECTE

1b RESTRICTIVE MARKINGS

NA

AUG 10 1988

3 DISTRIBUTION/AVAILABILITY OF REPORT

Distribution Unlimited

AD-A198 569

EDULE

NUMBER(S)

5 MONITORING ORGANIZATION REPORT NUMBER(S)

NA

6a NAME OF PERFORMING ORGANIZATION

University of Kansas

6b OFFICE SYMBOL  
(if applicable)

NA

7a NAME OF MONITORING ORGANIZATION

Office of Naval Research

6c ADDRESS (City, State, and ZIP Code)

Lawrence, KS 66045

7b ADDRESS (City, State, and ZIP Code)

800 N. Quincy St.  
Arlington, VA 22217-50008a NAME OF FUNDING/SPONSORING  
ORGANIZATION

Office of Naval Research

8b OFFICE SYMBOL  
(if applicable)

ONR

9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

N00014-86-K-0862

8c ADDRESS (City, State, and ZIP Code)

800 N. Quincy St.  
Arlington, VA 22217-5000

10 SOURCE OF FUNDING NUMBERS

PROGRAM  
ELEMENT NO

61153N

PROJECT  
NO

RR04106

TASK  
NO

NR 448016

WORK UNIT  
ACCESSION NO

11 TITLE (Include Security Classification)

Development of Synthetic Catalysts for Peptide Bond Cleavage  
Synthesis and Complete Kinetic Analysis of Compounds 6A, 7A, 8A

12 PERSONAL AUTHOR(S)

Mertes, Kristin Bowman, P.I., Mertes, Mathias Peter, Co-P.I.

13a TYPE OF REPORT

Annual

13b TIME COVERED

FROM 8/87 TO 8/88

14 DATE OF REPORT (Year, Month, Day)

1988, August 6

15 PAGE COUNT

4

16 SUPPLEMENTARY NOTATION

17 COSATI CODES

FIELD

GROUP

SUB-GROUP

18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Enzyme mimics; Supramolecular; Carboxypeptidase A;  
Polyammonium macrocycles

19 ABSTRACT (Continue on reverse if necessary and identify by block number)

Synthetic mimics for carboxypeptidase A will be synthesized and the structural and chemical factors responsible for catalytic peptidase activity will be probed. Ditopic macrocyclic receptors have been designed which incorporate the salient features of the enzyme analog, namely high affinity complex formation, general base and general acid catalysis, and covalent catalysis. Once synthesized the resulting macrocycle-metal ion complexes should non-specifically promote the hydrolysis of C-terminal peptide bonds. The initial macrocycles will have several types of coordination sites: nitrogen-containing heterocycles, ammonium and ether oxygens. One side of the ditopic receptor will preferentially bind zinc(II) ion, the other the peptide substrate.

## DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

20 DISTRIBUTION/AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS

21 ABSTRACT SECURITY CLASSIFICATION

(U)

22a NAME OF RESPONSIBLE INDIVIDUAL

Dr. M. Marron

22b TELEPHONE (Include Area Code)

202/696-4760

22c OFFICE SYMBOL

ONR

Date: August 6, 1988

ANNUAL PROGRESS REPORT ON CONTRACT N00014-86-K-0862

TASK NO. 44g016

PRINCIPAL INVESTIGATOR: Kristin Bowman Mertes

CO-PRINCIPAL INVESTIGATOR: Mathias P. Mertes

CONTRACT TITLE: Development of Synthetic Catalysts for Peptide Bond  
Cleavage: Synthesis and Complete Kinetic Analysis of  
Compounds 6A, 7A, 8A


START DATE: 6 August 1986

RESEARCH OBJECTIVE: To synthesize three macrocyclic ditopic receptors as mimics for carboxypeptidase A, i.e. as hydrolytic catalysts for ester and amide bonds alpha to a carboxylate group.


PROGRESS (YEAR 2): The report for the first year concentrated on the most optimal synthetic route for the preparation of the azoxychain representing the 'eastern' portion of the three target compounds shown in Figure 1. One of the major problems encountered was a polymerization reaction where we could not adequately control the number of units of the tosylaziridine that added to the azoxychain (8, Scheme 1, 87 report). This was complicated by the finding that the three major isomers were not easily separated. A second problem was the cyclization of the activated derivative (14, Scheme 1, 87 report) to give the morpholine.

These problems have now been solved using a convergent sequence as shown in Scheme 1. In this method the two halves, 3 and 6 are combined to give 7. This avoids the difficulties in extension of 5 which could not be controlled. While the overall yield is not good, simple starting materials combined with easier purification of the intermediates have afforded a reasonable amount of 8.

Reaction of 10 (87 report) with 9 afforded a reasonable yield of the protected macrocycle 11. Deprotection of 11 was a critical problem for several months until precise reagent quantities and temperature control were found. The problem encountered was that the side chain was being eliminated as acrylonitrile (Retro-Michael reaction). The amide 12 was characterized by <sup>1</sup>H and <sup>13</sup>C NMR, mass spec, and elemental analysis. Acid catalyzed hydrolysis of the amide to the acid 6A gave a product that had the correct NMR. However, we are concerned about the fact that we cannot get the molecular ion in the mass spec; we only see the M<sup>+</sup> + 1 - H<sub>2</sub>O. While the fragmentation pattern supports the structure the mass spec peak could also arise from a cyclic amide formed in the conversion of 12. IR data shows a carbonyl shift from 1670 (amide) to 1720 (COOH?). Further work will be done to assure that the acid 6A is the product and not a lactam by cyclization.




Scheme 1. Convergent route for the preparation of 9, the 'eastern' half of 6A, 7A, 8A.



Scheme 2. Synthesis of 6A.


Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

The 'western' half of compound **7A** was prepared in low overall yield as shown in Scheme 3 starting with **13**.



Scheme 3. Synthesis of the bipyridyl derivative **16**, the 'western' portion of **7A**.

Using the methods worked out for the first target compound, **7A** was prepared by reaction of **16** with the activated derivative **9** (Scheme 4). Again, controlled temperature and reaction monitoring were essential to minimize polymer formation. Conversion of **17** to the amide **18** proceeded as before. However, the amide hydrolysis to **7A** again afforded a product without the mass spectral molecular ion, only the dehydration product. Steps will be taken to verify that the structure is **7A**, not a lactam.



Scheme 4. Synthesis of **7A**

WORK PLAN (YEAR 3)

1. Structures 6A and 7A will be established.
2. Compound 8A will be synthesized using the methods worked out during the first two years.
3. Metal ( $\text{Zn}^{2+}$ ) complexation with these macrocycles will be studied.
4. The assay for peptidase and esterase activity using the macrocycles and their metal complexes will be explored using HPLC to improve the literature procedures. The amide precursors will also be studied in these assays to differentiate nucleophilic catalysis that could arise in the acids 6-8A.